

Hierarchical Investigation of Socioeconomic Drivers of Decadal Scale Land-Cover Changes in the Upper Midwest

PI: **Daniel G. Brown, University of Michigan**

Co-I: **Bryan C. Pijanowski, Michigan State University**

Co-I: **J. Michael Vasievich, USDA Forest Service**

URL: *www.umich.edu/~danbrown/research/nasa/nasa.html*



Introduction


RESEARCH QUESTIONS

- How can we observe and model changes in land use and land cover and their spatial patterns on a regional scale?
- What regional and local factors are driving spatial pattern of land use change?
- How is land use change affecting the amount and quality of forest cover in the region?
 - has implications for ecosystem health, forest resource production, carbon sequestration

APPROACH

- A combination of satellite imagery & aerial photography for observation and GIS-based spatial and statistical modeling.


ILLUSTRATION OF THE DATA. Our project created three-date maps of forest cover regionally, using the NALC data set, and land use within 136 sample sites (2500 ha), using aerial photography and digitized parcel maps.



Results

- Average increase in forest cover for our 136 sample sites was 17 % per decade, driven by ag. abandonment at an average rate of about 10 % by area per decade. Development, affecting an average of 30 % by area per decade, tended to increase forest clearing and dampen regrowth.
- We developed methods to estimate uncertainty in metrics of changing forest amount and pattern; global metrics tend to be more uncertain.
- We developed a two-step modeling process: (1) land use change as a function of exogenous drivers & (2) forest cover change as a stochastic simulation conditioned on land use change.
- Future steps include adding temporal depth to our data base, to improve our description of process, and incrementally refining the models.

CHANGE OCCURRING AT ONE SAMPLE SITE. Three land use classes: developed (red), undeveloped (green), and agriculture (yellow). Two cover classes: forest (green), not forest (gray).



Conclusions

- Forest cover increase in the Upper Midwest was real and should result in increased carbon sequestration, because it was replacing, mostly, pastures and croplands.
- The increase in forest cover was due, largely, to the abandonment of marginal agricultural lands as the farm economy has declined, a decline that could slow as marginal farmlands are removed from production and agriculture becomes more specialized. A significant lag effect was observed between abandonment and regrowth.
- A countervailing trend was extensive dispersed rural development in the region. This development was driven by demand for recreation and retirement destinations as well as for seasonal homes.
- To effectively map and model land use and land cover change in this region we required an approach that treated land use change, at the parcel level, and cover change, at the pixel level, as separate but linked processes.

PRESENTATIONS

- Brown, D.G. and Duh, J-D. 1998. Application of North American Landscape Characterization (NALC) Data for Studying Regional-Scale Landscape Pattern and Change. 13th Annual Meeting US Intl. Assoc. for Landscape Ecology, East Lansing, MI, March 20, 1998.
- Brown, D.G., Pijanowski, B.C., and Vasievich, J.M. 1998. Relationships Between Parcel Fragmentation and Changes in Land Use and Land Cover Patterns in the Upper Midwest USA.” Earth’s Changing Land, GCTE-LUCC Open Science Conference, Barcelona, Spain, March 16, 1998.
- Brown, D.G., Duh, J.D., 1999. "Estimating error in fragmentation metric values obtained using remote sensing for change analysis. 5th World Congress, Intl. Assoc. for Landscape Ecology, Snowmass, CO, July 1999.
- Brown, D.G. 2000. Land-Use to Land-Cover Change in the Upper Midwest, USA: A Missing Link. GIS and Environmental Modelling 4, Banff, Alberta, Canada, September, 2000.

PUBLICATIONS

- Drzyzga, S.A., Brown, D.G. 1998. Land Parcelization and Forest Fragmentation in Three Forested Counties in Michigan. *Proceedings*, Society of American Foresters 1998 National Convention, Traverse City, MI: 129-135. Brown, D.G., Duh, J.D., and Drzyzga, S. 2000. Estimating error in an analysis of forest fragmentation change using North American Landscape Characterization (NALC) Data. *Remote Sensing of Environment*, 71:106-117.
- Brown, D.G., Jacquez, G.M., Duh, J.D., and Maruca, S. In Press. Accuracy of remotely sensed estimates of landscape change using patch- and boundary-based pattern statistics. In: Huevelink, G. et al., eds. *Spatial Accuracy Assessment in Natural Resources and Environmental Sciences*, Delft University Press.
- Brown, D.G. In Press. Characterizing the human imprint on landscapes for ecological assessment. In Bourgeron, P. and Jensen, M., Eds. *An Integrated Ecological Assessment Protocol*. New York: Springer.
- Drzyzga, S.A. Pending Revision. Land Ownership Parcelization and Forest Fragmentation in Three Forested Counties in Northern Lower Michigan. Master's Thesis. East Lansing: Dept of Geography, Mich. State U.
- Brown, D.G., Pijanowski, B.C., and Duh, J.D. In Review. Modeling the relationships between land-use and land-cover on private lands in the Upper Midwest, USA. Submitted *Journal of Environmental Management*.
- Pijanowski, B.C., Brown, D.G., Manik, G., Shellito, B.A. In Preparation. Using neural nets and GIS to forecast land use changes: A land transformation model.
- Brown, D.G., Pijanowski, B.C., Savage, S.E. In Preparation. Effects of land-use change on forest-cover in the Upper Midwest at the parcel scale.